

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

CTP, Zewail City for Science and Technology, Egypt
April 19, 2012

Mary Bishai (on behalf of the Daya Bay Collaboration)





Outline

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

1 Introduction

- History of Neutrinos
- Neutrino Mixing

2 The Daya Bay Experiment

- Introduction
- Experimental Apparatus
- Signal Selection
- Calibration
- Backgrounds
- Results

3 Summary

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction

History of
Neutrinos
Neutrino Mixing


The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results


Summary

INTRODUCTION TO EXPERIMENTAL NEUTRINO PHYSICS

Neutrino Conception



Before 1930's: beta decay spectrum continuous - is this energy non-conservation?



The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction
History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Neutrino Conception

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Dec 1930: Wolfgang Pauli's letter to physicists at a workshop in Tübingen:



Wolfgang Pauli

Dear Radioactive Ladies and Gentlemen,

....., I have hit upon a desperate remedy to save the "exchange theorem" of statistics and the law of conservation of energy. Namely, the possibility that there could exist in the nuclei electrically neutral particles, that I wish to call neutrons.... The mass of the neutrons should be of the same order of magnitude as the electron mass and in any event not larger than 0.01 proton masses. The continuous beta spectrum would then become understandable by the assumption that in beta decay a neutron is emitted in addition to the electron such that the sum of the energies of the neutron and the electron is constant.....

Unfortunately, I cannot appear in Tübingen personally since I am indispensable here in Zurich because of a ball on the night of 6/7 December. With my best regards to you, and also to Mr Back.

Your humble servant

. W. Pauli

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction

History of
Neutrinos

Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus

Signal Selection
Calibration
Backgrounds
Results

Summary

1932: **James Chadwick** discovers the neutron,
 $\text{mass}_{\text{neutron}} = 1.0014 \times \text{mass}_{\text{proton}}$ - its too heavy -
cant be Pauli's particle



James Chadwick

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)


Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Solvay Conference, Bruxelles 1933: **Enrico Fermi**
proposes to name Pauli's particle the "**neutrino**".



Enrico Fermi

The Theory of Weak Interactions

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction
History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment


Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary


≥ 1933: Fermi Glashow/Salam/Weinberg (1968) build the theory of **weak interactions and beta decay**

Charged current interactions

Decay of neutron




Neutrino interacts with neutron



Neutral current interactions

n or p interacts with neutrino or antineutrino



Finding Neutrinos...

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary


1950's: Fred Reines at Los Alamos and Clyde Cowan use the Hanford nuclear reactor (1953) and the new Savannah River nuclear reactor (1955) to find neutrinos. A detector filled with **water with CdCl_2 in solution** was located 11 meters from the reactor center and 12 meters underground.

The detection sequence was as follows:

- 1 $\bar{\nu}_e + p \rightarrow n + e^+$

- 2 $e^+ + e^- \rightarrow \gamma\gamma$

- 3 $n + {}^{108}\text{Cd} \rightarrow {}^{109}\text{Cd}^* \rightarrow {}^{109}\text{Cd} + \gamma$
($\tau = 5\mu\text{s}$).



Neutrinos first detected using a nuclear reactor!

ν : A Truly Elusive Particle!

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Reines and Cowan were the first to estimate the interaction strength of neutrinos.

The cross-section is $\sigma \sim 10^{-43} \text{cm}^2$ per nucleon (p,n).

$$\nu \text{ mean free path} = \frac{\text{Mass of the proton}}{\sigma \times \text{density}}$$

$$= \frac{1.67 \times 10^{-24} \text{g}}{10^{-43} \text{cm}^2 \times 11.4 \text{g/cm}^3} \approx 1.5 \times 10^{16} \text{m} = \mathbf{1.6 \text{ LIGHT YEARS}}$$

A proton has a mean free path of 10cm in lead

Neutrino detectors have to be MASSIVE


Neutrinos have Flavors



1962: Leon Lederman, Melvin Schwartz and Jack Steinberger use BNL's Alternating Gradient Synchrotron (AGS) to produce a beam of neutrinos using the decay $\pi \rightarrow \mu \nu_x$



The AGS



Making ν 's

Result: 40 neutrino interactions recorded in the detector, 6 of the resultant particles were identified as background and 34 identified as $\mu \Rightarrow \nu_x = \nu_\mu$

The first accelerator neutrino experiment was at the AGS.

1988 Nobel prize


Number of Neutrino Flavors: Particle Colliders

1980's - 90's: The number of neutrino types is precisely determined from studies of Z^0 boson properties produced in e^+e^- colliders.

The LEP e^+e^- collider at CERN, Switzerland



The 27km LEP ring was reused to
build the Large Hadron Collider



$$N_\nu = 2.984 \pm 0.008$$

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)


Introduction

History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



Sources of Neutrinos

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction

History of Neutrinos

Neutrino Mixing

The Daya Bay Experiment

Introduction

Experimental Apparatus

Signal Selection


Calibration

Backgrounds

Results


Summary

Big Bang




10^{-4} eV
 $300/\text{cm}^3$

Reactors




few MeV
 $10^{21}/\text{GW}_{\text{th}}/\text{s}$

Sun




0.1-14 MeV
 $10^{10}/\text{cm}^2/\text{s}$

SuperNova



~ 10 MeV
 $10^9/\text{cm}^2/\text{s}$

Atmosphere



~ 1 GeV
few/ cm^2/s

Accelerators



1-20 GeV
 $10^5/\text{cm}^2/\text{s}$ (at 1km)

Extragalactic



TeV-PeV
varies

beta-decay limits)

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction
History of Neutrinos
Neutrino Mixing


The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary


$$\Omega_i \equiv \rho_i / \rho_{\text{CRITICAL}}$$

$$\Omega_{\text{TOTAL}} = 1$$



Heavy Elements:

$$\Omega=0.0003$$




Neutrinos (ν):

$$\Omega=0.0047$$




Stars:

$$\Omega=0.005$$




Free H & He:

$$\Omega=0.04$$



Cold Dark Matter:

$$\Omega=0.25$$



Dark Energy (Λ):

$$\Omega=0.70$$

Neutrino oscillations

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction

History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

1957,1967: **B. Pontecorvo** proposes that neutrinos could oscillate:

$$\begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}$$

$$\nu_a(t) = \cos(\theta)\nu_1(t) + \sin(\theta)\nu_2(t)$$


$$\begin{aligned} P(\nu_a \rightarrow \nu_b) &= |\langle \nu_b | \nu_a(t) \rangle|^2 \\ &= \sin^2(\theta) \cos^2(\theta) |e^{-iE_2 t} - e^{-iE_1 t}|^2 \end{aligned}$$

$$P(\nu_a \rightarrow \nu_b) = \sin^2 2\theta \sin^2 \frac{1.27 \Delta m_{21}^2 L}{E}$$

where $\Delta m_{21}^2 = (m_2^2 - m_1^2)$ in eV^2 ,
 L (km) and E (GeV).

Observation of oscillations

implies non-zero mass eigenstates



The Homestake Experiment

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

1967: Ray Davis from BNL installs a large detector, containing 615 tons of tetrachloroethylene (cleaning fluid), 1.6km underground in Homestake mine, SD.


1 $\nu_e^{\text{sun}} + {}^{37}\text{Cl} \rightarrow e^- + {}^{37}\text{Ar}$, $\tau({}^{37}\text{Ar}) = 35$ days.

2 Number of Ar atoms \approx number of ν_e^{sun} interactions.



Ray Davis

Results: 1969 - 1993 Measured 2.5 ± 0.2 SNU (1 SNU = 1 neutrino interaction per second for 10^{36} target atoms) while theory predicts 8 SNU:



This is a ν_e^{sun} deficit of 69%.

Solar ν_e disappearance \Rightarrow

first experimental hint of oscillations (2002 Nobel prize)

SNO Experiment: Solar ν Measurements

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction
History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

$1 \leftrightarrow 2$ mixing

2001-02: Sudbury Neutrino Observatory. Water Čerenkov detector with 1 kT heavy water (**0.5 B\$ worth on loan from Atomic Energy of Canada Ltd.**) located 2Km below ground in INCO's Creighton nickel mine near Sudbury, Ontario. Can detect the following ν^{sun} interactions:

- 1) $\nu_e + d \rightarrow e^- + p + p$ (CC).
- 2) $\nu_x + d \rightarrow p + n + \nu_x$ (NC).
- 3) $\nu_x + e^- \rightarrow e^- + \nu_x$ (ES).


SNO measured:

$$\phi_{\text{SNO}}^{\text{CC}}(\nu_e) = 1.75 \pm 0.07(\text{stat})_{-0.11}^{+0.12}(\text{sys.}) \pm 0.05(\text{theor}) \times 10^6 \text{cm}^{-2} \text{s}^{-1}$$

$$\phi_{\text{SNO}}^{\text{ES}}(\nu_x) = 2.39 \pm 0.34(\text{stat})_{-0.14}^{+0.16}(\text{sys.}) \pm \times 10^6 \text{cm}^{-2} \text{s}^{-1}$$

$$\phi_{\text{SNO}}^{\text{NC}}(\nu_x) = 5.09 \pm 0.44(\text{stat})_{-0.43}^{+0.46}(\text{sys.}) \pm \times 10^6 \text{cm}^{-2} \text{s}^{-1}$$

All the solar ν 's are there but ν_e appears as ν_x !



The Implications of 3-Neutrino Mixing

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction

History of Neutrinos

Neutrino Mixing

The Daya Bay Experiment

Introduction

Experimental Apparatus

Signal Selection

Calibration

Backgrounds

Results

Summary


$$U_{\text{PMNS}} = \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix}}_{\nu_\mu \text{ disappearance}} \underbrace{\begin{pmatrix} c_{13} & 0 & e^{i\delta_{\text{CP}}} s_{13} \\ 0 & 1 & 0 \\ -e^{i\delta_{\text{CP}}} s_{13} & 0 & c_{13} \end{pmatrix}}_{\nu_\mu \rightarrow \nu_e, \text{ reactor } \bar{\nu}_e \text{ disappear}} \underbrace{\begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\text{solar } \nu_e, \bar{\nu}_e \text{ disappear}}$$

where $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$.

$\sin^2 \theta_{13}$: Amount of ν_e in ν_3

$\tan^2 \theta_{23}$: Ratio of $\frac{\nu_\mu}{\nu_\tau}$ in ν_3

$\tan^2 \theta_{12}$: $\frac{\text{Amount of } \nu_e \text{ in } \nu_2}{\text{Amount of } \nu_e \text{ in } \nu_1}$



Normal

Inverted

There are 3 quantum states mixing \Rightarrow there is an overall phase: δ_{CP} .

If $\delta_{\text{CP}} \neq 0$ or π , charge-parity (CP) is violated and there is a $\nu/\bar{\nu}$ asymmetry.

Could this explain the origin of matter?

What is the value of δ_{CP} , $\text{sign}(\Delta m_{31}^2)$?




KamLAND: Reactor $\bar{\nu}_e$ Measurements

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

- Introduction
- History of Neutrinos
- Neutrino Mixing**
- Neutrino Masses
- Neutrino Oscillations
- Neutrinoless Double Beta Decay
- Neutrino Physics at the LHC
- Neutrino Physics at the FCC

1 \leftrightarrow 2 mixing



⇐ **Clear wiggles!**

$$\Delta m_{21}^2 = 7.59^{+0.21}_{-0.21} \times 10^{-5} \text{ eV}^2$$

$$\tan^2 \theta_{12} = 0.47^{+0.06}_{-0.05}$$


Measuring neutrino mixing - ν_e oscillations

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

- Introduction
- History of Neutrinos
- Neutrino Mixing

Solar ν_e disappearance constrained $1 \rightarrow 2$ mixing. Precision from reactor $\bar{\nu}_e$ experiments :



The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction

History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Observation of $\bar{\nu}_e$ Disappearance at Daya Bay

Short Baseline Reactor $\bar{\nu}_e$ oscillations

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)


Introduction
History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

$$P(\nu_e \rightarrow \nu_e) = 1 - \sin^2 2\theta_{13} \sin^2(1.27 \Delta m_{31}^2 L/E) - \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2(1.27 \Delta m_{21}^2 L/E)$$



**Unambiguous measurement
of $\sin^2 2\theta_{13}$**

Getting to $\sin^2 2\theta_{13} < 0.01$

Lots of statistics: -Powerful nuclear reactors + more massive detectors

Suppress cosmic backgrounds:

-Increase overburden = go deeper underground.

Reduce systematic uncertainties:

-Deploy near detectors as close as possible to reactor to minimize reactor flux uncertainties.

-Use "identical" N/F detectors to reduce near/far detector uncertainties.

-Calibration, calibration, calibration...

Detecting $\bar{\nu}_e$ using GD-loaded LS.

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)



Introduction
History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

The active target in each detector module is liquid scintillator loaded with 0.1% Gd



The detection sequence is as follows: $\bar{\nu}_e + p \rightarrow n + e^+$ THEN

$e^+ + e^- \rightarrow \gamma\gamma$ (2X 0.511 MeV + T_{e^+} , prompt)

$n + p \rightarrow D + \gamma$ (2.2 MeV, $\tau \sim 180\mu s$, $\sigma = 0.3b$). OR

$n + Gd \rightarrow Gd^* \rightarrow Gd + \gamma's$ (8 MeV, $\tau \sim 28\mu s$, $\sigma = 5 \times 10^4 b$).

\Rightarrow delayed co-incidence of e^+ conversion and n-capture (> 6 MeV)

with a specific energy signature

The Daya Bay Reactor Complex

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction

History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



Reactor Specs:

Located 55km north-east of Hong Kong.

2 cores at Daya Bay site + 2 cores at Ling Ao site = 11.6 GW_{th}

2 more cores at Ling Ao II site = 17.4 GW_{th} ⇒ top five worldwide

1 GW_{th} = $2 \times 10^{20} \bar{\nu}_e$ /second

Deploy multiple near and far detectors

Reactor power uncertainties < 0.1%

The Daya Bay Collaboration

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Political Map of the World, June 1999



The Daya Bay Experimental Apparatus

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications



Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



- Multiple “identical” detectors at each site.
- Manual and multiple automated calibration systems per detector.
- Thick water shield/muon veto to reduce cosmogenic and radiation bkgds.

	DYB	LA	Far
Event rates/20T/day	840	740	90

Daya Bay Measurement Technique

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai
(on behalf of the Daya Bay Collaboration)

Introduction


History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction

Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



$$\frac{N_f}{N_n} = \left(\frac{N_{p,f}}{N_{p,n}} \right) \left(\frac{L_n}{L_f} \right)^2 \left(\frac{\epsilon_f}{\epsilon_n} \right) \left[\frac{P_{\text{sur}}(E, L_f)}{P_{\text{sur}}(E, L_n)} \right]$$

Far/Near
Neutrino Ratio

Detector
Target Mass

Distance
from
Reactor

Detector
Efficiency

Survival Probability
(θ_{13})

Daya Bay Measurement Goals

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction

History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Source of uncertainty		Chooz (<i>absolute</i>)	Daya Bay (<i>relative</i>)	Strategy
# protons	H/C ratio	0.8	< 0.1	Fill in pairs/calib
	Mass	-	< 0.3	Load cells and mass flowmeters
Detector Efficiency	Energy cuts	0.8	0.2	lower threshold/calib
	Position cuts	0.32	0.0	3-zone
	Time cuts	0.4	0.1	Common clock ~ 10ns
	H/Gd ratio	1.0	0.1	fill in pairs/calib
	n multiplicity	0.5	0.05	Deeper/muon veto
	Trigger	0	0.01	Redundant triggers
	Live time	0	< 0.01	Common GPS clock
Total detector-related uncertainty		1.7%	0.38%	

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction


History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus


Signal Selection
Calibration
Backgrounds
Results

Summary





Daya Bay Liquid Scintillator Cocktail

- LAB + Gd (0.1%) + PPO (3 g/L) + bis-MSB (15 mg/L)
- more than 3 years R&D (BNL & IHEP)
- Multi-stage purifications on optical improvement and U/Th removal
- 185-ton Gd-LS + 196-ton LS production



Load cells measure 20 ton target mass to 3kg (0.015%)



Anti-Neutrino Detector Assembly

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications






Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing



The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results



Summary



EH1 (Daya Bay) 15 Aug 2011



EH2 (Ling Ao I, II) 5 Nov 2011



EH3 (far site)

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)



Introduction

History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary




Energy Spectrum of All Events

Using a ^{60}Co source (2.506 MeV) at center:

~ 163 photo-electrons/MeV

All triggered events




Energy Spectrum of All Events

Using a ^{60}Co source (2.506 MeV) at center:

~ 163 photo-electrons/MeV

Veto muons: water shield $\mu > 0.6\text{ms}$, AD $\mu (> 20 \text{ MeV}, > 1 \text{ ms})$, AD shower $\mu (> 2.5 \text{ GeV}, > 1 \text{ s})$



Identifying the Inverse Beta Decay Signal

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Selection criteria:



Muon veto

Prompt positron: $0.7 < E_{\text{prompt}} < 12 \text{ MeV}$

Delayed neutron: $6 < E_{\text{delayed}} < 12 \text{ MeV}$


Capture time: $1 < t_{\text{delayed}} - t_{\text{prompt}} < 200 \mu\text{s}$

Multiplicity: $< 0.7 \text{ MeV}$ in the time range
($t_p - 200 \mu\text{s}, t_d + 200 \mu\text{s}$).





I.B.D Rates and Spectra in EH1

Positron ($\bar{\nu}_e$) signal



Neutron capture signal



The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction


History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Automated Calibration Units (ACUs)



The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction

History of Neutrinos
Neutrino Mixing


The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

LEDs and 2" Calibration PMTs

- LEDs along three deployment axes.
- 3 fixed LEDs on detector inner wall.
- 2" PMTs on the top and bottom of detector.
- Monitor optical properties
- PMT calibration



The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction


History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment


Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Single pe peak




PMT gain stability




Gamma sources: Ge-68 & Co-60


- Ge-68 (Rate: 100 Bq, $T_{1/2}$: 270 days)



- Positron threshold
- Relative PMT detecting efficiency
- Co-60 (Rate: 150 Bq, $T_{1/2}$: 1925 days)




- Energy calibration
- Monitor light yield/attenuation



Neutron source: AmC
(Next slide)

Neutron source: Am-C



$^{241}\text{Am}-^{13}\text{C}$ (Rate: 0.5 Hz)

- $^{13}\text{C} (\alpha, n) ^{16}\text{O}$
- Au foil attenuate α to < 4.5 MeV, hence suppressing 6.13 MeV gamma from excited state of ^{16}O .
- Neutron energy scale



Combined Source Calibration Spectrum Am-¹³C/⁶⁰Co EH1

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications


Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



Energy Calibration - Position Non-Uniformities

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)


Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment


Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

With different sources



EH1 and 2



Energy Calibration - Resolution

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)


Introduction

History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment


Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



Background Accidentals

Two uncorrelated events mimic the anti-neutrino (IBD) signals



Rate and spectra can be accurately predicted from the singles data

Most delayed-like single events are from beta-decays of long-live muon spallation isotopes

B/S ratio is ~4.5% at Far site, ~1.4% at Near site

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction

History of Neutrinos
Neutrino Mixing

The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary

Background Muon Spallation

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction

History of Neutrinos
Neutrino Mixing


The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary


β -n decay:

- Prompt: β -decay
- Delayed: neutron capture





$${}^9\text{Li}: \tau_{1/2} = 178 \text{ ms}, Q = 13.6 \text{ MeV}$$

$${}^8\text{He}: \tau_{1/2} = 119 \text{ ms}, Q = 10.6 \text{ MeV}$$



B/S: 0.2%



B/S: 0.06%

Neutrino Data in 3 Experimental Halls

The Measurement of the Neutrino Mixing Angle, θ_{13} , and its Implications

Mary Bishai (on behalf of the Daya Bay Collaboration)

Introduction
History of Neutrinos
Neutrino Mixing


The Daya Bay Experiment

Introduction
Experimental Apparatus
Signal Selection
Calibration
Backgrounds
Results


Summary

	AD1	AD2	AD3	AD4	AD5	AD6
IBD candidates	28935	28975	22466	3528	3436	3452
DAQ live time (days)	49.5530		49.4971	48.9473		
$\epsilon_{\mu} \cdot \epsilon_m$	0.8019	0.7989	0.8363	0.9547	0.9543	0.9538
Accidentals (per day)	9.82±0.06	9.88±0.06	7.67±0.05	3.29 ±0.03	3.33 ± 0.03	3.12 ±0.03
Fast-neutron (per day)	0.84±0.28	0.84±0.28	0.74±0.44	0.04±0.04	0.04±0.04	0.04±0.04
$^9\text{Li}/^8\text{He}$ (per AD per day)	3.1±1.6		1.8±1.1	0.16±0.11		
Am-C correlated (per AD per day)	0.2±0.2					
$^{13}\text{C}(\alpha, n)^{16}\text{O}$ background (per day)	0.04±0.02	0.04±0.02	0.035±0.02	0.03±0.02	0.03±0.02	0.03±0.02
IBD rate (per day)	714.17±4.58	717.86± 4.60	532.29±3.82	71.78 ± 1.29	69.80±1.28	70.39±1.28

High statistics anti-neutrino spectra



No-oscillation predictions are obtained using reactor flux analysis and detector simulations and compared to the observed I.B.D. rate:



Detector			
	Efficiency	Correlated	Uncorrelated
Target Protons		0.47%	0.03%
Flasher cut	99.98%	0.01%	0.01%
Delayed energy cut	90.9%	0.6%	0.12%
Prompt energy cut	99.88%	0.10%	0.01%
Multiplicity cut		0.02%	<0.01%
Capture time cut	98.6%	0.12%	0.01%
Gd capture ratio	83.8%	0.8%	<0.1%
Spill-in	105.0%	1.5%	0.02%
Livetime	100.0%	0.002%	<0.01%
Combined	78.8%	1.9%	0.2%

Reactor			
Correlated		Uncorrelated	
Energy/fission	0.2%	Power	0.5%
IBD reaction/fission	3%	Fission fraction	0.6%
		Spent fuel	0.3%
Combined	3%	Combined	0.8%

Measurement of $\sin^2 2\theta_{13}$: March 8, 2012

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)



Introduction

History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



Rate only: $\sin^2 2\theta_{13} = 0.092 \pm 0.016(\text{stat}) \pm 0.005(\text{syst})$

- The Daya Bay experiment has measured a non-zero value for the neutrino mixing angle θ_{13} with a significance of 5.2σ . This is the first observation of a non-zero θ_{13} .
- A rate only analysis finds:
Rate only: $\sin^2 2\theta_{13} = 0.092 \pm 0.016(\text{stat}) \pm 0.005(\text{syst})$
- On April 12, 2012 the RENO experiment confirmed the Daya Bay result and measured:
Rate only: $\sin^2 2\theta_{13} = 0.113 \pm 0.013(\text{stat}) \pm 0.019(\text{syst})$
- The stage is now set for measurement of the CP-violating phase and the hunt for CP violation in the neutrino sector using $\nu_{\mu} \rightarrow \nu_e$ oscillations with the next generation of accelerator long baseline neutrino experiments.

The
Measurement
of the
Neutrino
Mixing Angle,
 θ_{13} , and its
Implications

Mary Bishai
(on behalf of
the Daya Bay
Collaboration)

Introduction
History of
Neutrinos
Neutrino Mixing

The Daya Bay
Experiment

Introduction
Experimental
Apparatus
Signal Selection
Calibration
Backgrounds
Results

Summary



THANK YOU